



Contents lists available at ScienceDirect

Maturitas

journal homepage: www.elsevier.com/locate/maturitas

Editorial

The interconnection between Covid-19, sarcopenia and lifestyle

ARTICLE INFO

Keywords

Sarcopenia

Lifestyle

Covid-19

In March 2020, Covid-19 was declared a global pandemic by the World Health Organization. Infection with Sars-Cov-2, the virus responsible for Covid-19, can have a wide range of consequences, the severity of which depends on the individual [1]. Moreover, this vulnerability is increased by the presence of comorbidities such as chronic respiratory disease, diabetes, hypertension and other pathologies affecting immunity [2]. Also, musculoskeletal disorders, and particularly sarcopenia, have recently been suggested to be either a consequence of [3] or a risk factor for [4] Covid-19. For example, studies have reported muscle damage in patients hospitalized for Covid-19 and greater loss of muscle function in older women living independently during the pandemic [5,6]. Some studies, but not all, have also highlighted that low skeletal muscle mass and reduced muscle strength are associated with higher severity of Covid-19 [7–9]. Very recently, histopathological changes in muscle, such as muscle atrophy, have also been identified in patients complaining of long-term Covid-19 symptoms, including fatigue and weakness [10]. In addition, results from a meta-analysis highlighted a twofold greater risk of getting severe complications and of mortality due to Covid-19 for sarcopenic people compared with non-sarcopenic ones [4].

Three potentially modifiable factors appear to be associated with both sarcopenia and Covid-19 and can be considered a target for prevention: high level of systemic inflammation, low levels of physical activity and poor nutrition:

- *High level of systemic inflammation:* Sarcopenia can potentially induce a higher level of inflammation, which can be observed with trough biomarkers such as interleukin-15 (IL-15), interleukin-6 (IL-6) and C-reactive protein (CRP) [11,12]. In parallel, in severe forms of Covid-19 disease, a high level of biomarkers of inflammation (i.e. IL-6 and IL-10) have been found [13] and can potentially increase the risk of acute sarcopenia [14].
- *Low physical activity:* The model of “catabolic crisis”, which proposes that sarcopenia is not only like a progressive process but also aggravated by acute periods of inactivity, was meaningful during the Covid-19 crisis [15]. Indeed, quarantine, lockdown restriction and

hospitalization were among the many conditions that affected lifestyle behaviours and consequently might have increased the risk of developing sarcopenia. In this context, a systematic review showed an overall global trend of decreased physical activity in older adults due to lockdown restrictions [16]. The authors of this review noted that in addition to induced lower muscle mass and strength, which is directly associated with sarcopenia, lower levels of physical activity can lead to other non-communicable diseases that increase vulnerability to Covid-19 [16].

- *Poor nutrition:* Some harmful eating behaviours have been reported during Covid-19 quarantine, such as skipping warm meals, eating less than normal and having less appetite [6]. The impact of these changes was partially highlighted in a study in which 25 % of 1407 patients hospitalized for Covid-19 had significant weight loss before hospitalization [17]. In that study, this malnourishment could be partly explained by Covid-19-related symptoms like ageusia, change of taste, nausea, vomiting, and anorexia [17]. Those changes in eating habits may cause a decrease in protein intake. Indeed, insufficient intake can lead to a reduction in muscle protein synthesis which is an aggravating factor for sarcopenia [3]. In addition, the decrease in physical activity observed during the pandemic could indirectly lead to changes in other lifestyle behaviours, including those related to nutrition. Indeed, during the Covid-19 pandemic, people with lower physical activity levels increased snacking habits and alcohol consumption [6]. These eating habits may lead to overnutrition, as discussed in a Brazilian study conducted in older women during the pandemic that revealed an increase of body mass index and body weight but also an increased risk of sarcopenia [18].

The good news, at least for preventive actions for both Covid-19 and sarcopenia, is that inflammation, physical activity and nutrition are all potentially modifiable and interconnected factors. As specified by the International Clinical Practice Guidelines for Sarcopenia, physical activity and adapted diet are crucial in the management of sarcopenia [19]. For physical activity, to date, resistance-based training is still preferred for its benefits for muscle strength, physical performance and

<https://doi.org/10.1016/j.maturitas.2022.09.009>

Received 14 September 2022; Accepted 23 September 2022

Available online 6 October 2022

0378-5122/© 2022 Elsevier B.V. All rights reserved.

muscle hypertrophy [19]. Regarding nutrition, it is recommended that older adults are monitored to ensure energy balance as well as adequate intake of protein and vitamin D. Acting on these two together can generate a decrease in inflammation and further potentially reduce the risk of sarcopenia. Indeed, in their systematic review, Bautmans et al. have highlighted a substantial lower level of inflammatory markers such as IL-6 and CRP in older adults following a physical exercise program in comparison with those who have not [20]. In addition, a recent meta-analysis highlighted that healthy eating behaviours such as high intake of fresh fruits are correlated with lower level of inflammatory biomarkers [21].

In conclusion, even though most restrictive rules have been lifted in almost all countries, Covid-19 is still active. On 15 August 2022, for one year 375 million new cases and over 2 million deaths were reported on a global scale [22]. At present, older populations are still considered vulnerable to Covid-19. Sarcopenia appears to be a risk factor for as well as a consequence of Covid-19 and has repercussions in terms of severity and mortality. In line with what some authors have called “a wake-up call for lifestyle-related preventable conditions in older adults” [23], public health authorities and health professionals [24] should be aware of the connection between sarcopenia and Covid-19 and the factors aiming at reducing their burden.

Contributors

Céline Demonceau participated in data collection and analysis, and in drafting and editing of the paper.

Charlotte Beaudart participated in data collection and analysis, and in drafting and editing of the paper.

Jean-Yves Reginster participated in data collection and analysis.

Nicola Veronese participated in data collection and analysis.

Olivier Bruyère participated in data collection and analysis, and in drafting and editing of the paper.

All authors saw and approved the final version and no other person made a substantial contribution to the paper.

Funding

The authors received no funding from an external source for the preparation of this editorial.

Provenance and peer review

This article was commissioned and was not externally peer reviewed.

Declaration of competing interest

The authors declare that they have no competing interest.

References

- [1] R. Guerrina, B. Borisch, L.F. Callahan, J. Howick, J.Y. Reginster, A. Mobasher, Health and gender inequalities of the COVID-19 pandemic: adverse impacts on women's health, wealth and social welfare, *Front Glob Womens Health* 2 (2021), 670310.
- [2] M.S. Gold, D. Sehayek, S. Gabrielli, X. Zhang, C. McCusker, M. Ben-Shoshan, COVID-19 and comorbidities: a systematic review and meta-analysis, *Postgrad. Med.* 132 (8) (2020) 749–755.
- [3] R. Kirwan, D. McCullough, T. Butler, F. Perez de Heredia, I.G. Davies, C. Stewart, Sarcopenia during COVID-19 lockdown restrictions: long-term health effects of short-term muscle loss, *Geroscience* 42 (6) (2020) 1547–1578.
- [4] Y.M.T. Siahaan, V. Hartoyo, T.I. Hariyanto, A. Kurniawan, Coronavirus disease 2019 (Covid-19) outcomes in patients with sarcopenia: A meta-analysis and meta-regression, *Clin Nutr ESPEN* 48 (2022) 158–166.
- [5] L. Mao, H. Jin, M. Wang, Y. Hu, S. Chen, Q. He, J. Chang, C. Hong, Y. Zhou, D. Wang, X. Miao, Y. Li, B. Hu, Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in WuhanChina, *JAMA Neurol* 77 (6) (2020) 683–690.
- [6] M. Visser, L.A. Schaap, H.A.H. Wijnhoven, Self-reported impact of the COVID-19 pandemic on nutrition and physical activity behaviour in dutch older adults living independently, *Nutrients* 12 (12) (2020).
- [7] Y. Ma, M. He, L.S. Hou, S. Xu, Z.X. Huang, N. Zhao, Y. Kang, J.R. Yue, C. Wu, The role of SARC-F scale in predicting progression risk of COVID-19 in elderly patients: a prospective cohort study in Wuhan, *BMC Geriatr.* 21 (1) (2021) 355.
- [8] X. Yi, H. Liu, L. Zhu, D. Wang, F. Xie, L. Shi, J. Mei, X. Jiang, Q. Zeng, P. Hu, Y. Li, P. Pang, J. Liu, W. Peng, H.X. Bai, W. Liao, B.T. Chen, Myosteatosis predicting risk of transition to severe COVID-19 infection, *Clin. Nutr.* (2021), <https://doi.org/10.1016/j.clnu.2021.05.031>. Epub ahead of print.
- [9] L. Lengele, M. Locquet, M. Moutschen, C. Beaudart, J.F. Kaux, S. Gillain, J. Y. Reginster, O. Bruyere, Frailty but not sarcopenia nor malnutrition increases the risk of developing COVID-19 in older community-dwelling adults, *Aging Clin. Exp. Res.* 34 (1) (2022) 223–234.
- [10] E.K. Hejbol, T. Harbo, J. Agergaard, L.B. Madsen, T.H. Pedersen, L.J. Ostergaard, H. Andersen, H.D. Schroder, H. Tankisi, Myopathy as a cause of fatigue in long-term post-COVID-19 symptoms: evidence of skeletal muscle histopathology, *Eur. J. Neurol.* 29 (9) (2022) 2832–2841.
- [11] C. Nelke, R. Dziewas, J. Minnerup, S.G. Meuth, T. Ruck, Skeletal muscle as potential central link between sarcopenia and immune senescence, *EBioMedicine* 49 (2019) 381–388.
- [12] G. Bano, C. Trevisan, S. Carraro, M. Solmi, C. Luchini, B. Stubbs, E. Manzato, G. Sergi, N. Veronese, Inflammation and sarcopenia: a systematic review and meta-analysis, *Maturitas* 96 (2017) 10–15.
- [13] S.K. Dhar, V. K. S. Damodar, S. Gujar, M. Das, IL-6 and IL-10 as predictors of disease severity in COVID-19 patients: results from meta-analysis and regression, *Heliyon* 7 (2) (2021), e06155.
- [14] K. Piotrowicz, J. Gasowski, J.P. Michel, N. Veronese, Post-COVID-19 acute sarcopenia: physiopathology and management, *Aging Clin. Exp. Res.* 33 (10) (2021) 2887–2898.
- [15] K.L. English, D. Paddon-Jones, Protecting muscle mass and function in older adults during bed rest, *Curr. Opin. Clin. Nutr. Metab. Care* 13 (1) (2010) 34–39.
- [16] A.L. Elisabeth, S.B. Karlen, F. Magkos, The effect of COVID-19-related lockdowns on diet and physical activity in older adults: a systematic review, *Aging Dis.* 12 (8) (2021) 1935–1947.
- [17] N.J. Wierdsma, H.M. Kruizenga, L.A. Konings, D. Krebbers, J.R. Jorissen, M. I. Joosten, L.H. van Aken, F.M. Tan, A.A. van Bodegraven, M.R. Soeters, P.J. Weijs, Poor nutritional status, risk of sarcopenia and nutrition related complaints are prevalent in COVID-19 patients during and after hospital admission, *Clin. Nutr. ESPEN* 43 (2021) 369–376.
- [18] A.Q. da Rocha, P.C.B. Lobo, G.D. Pimentel, Muscle function loss and gain of body weight during the COVID-19 pandemic in elderly women: effects of one year of lockdown, *J. Nutr. Health Aging* 25 (8) (2021) 1028–1029.
- [19] E. Dent, J.E. Morley, A.J. Cruz-Jentoft, H. Arai, S.B. Kritchevsky, J. Guralnik, J. M. Bauer, M. Pahor, B.C. Clark, M. Cesari, J. Ruiz, C.C. Sieber, M. Aubertin-Leheudre, D.L. Waters, R. Visvanathan, F. Landi, D.T. Villareal, R. Fielding, C. W. Won, O. Theou, F.C. Martin, B. Dong, J. Woo, L. Flicker, L. Ferrucci, R. A. Merchant, L. Cao, T. Cederholm, S.M.L. Ribeiro, L. Rodriguez-Manas, S. D. Anker, J. Lundy, L.M. Gutierrez Robledo, I. Bautmans, I. Aprahamian, J. Schols, M. Izquierdo, B. Vellas, International clinical practice guidelines for sarcopenia (ICFSR): screening, diagnosis and management, *J. Nutr. Health Aging* 22 (10) (2018) 1148–1161.
- [20] I. Bautmans, L. Salimans, R. Njemini, I. Beyer, S. Lieten, K. Liberman, The effects of exercise interventions on the inflammatory profile of older adults: a systematic review of the recent literature, *Exp. Gerontol.* 146 (2021), 111236.
- [21] M.M. Norde, T.S. Collese, E. Giovannucci, M.M. Rogero, A posteriori dietary patterns and their association with systemic low-grade inflammation in adults: a systematic review and meta-analysis, *Nutr. Rev.* 79 (3) (2021) 331–350.
- [22] W.H. Organization, COVID-19 Weekly Epidemiological Update, 2022.
- [23] M.A. Lim, L. Smith, COVID-19 pandemic: a wake-up call for lifestyle-related preventable conditions in older adults, *Aging Clin. Exp. Res.* 33 (8) (2021) 2367–2368.
- [24] Z. Altin, F. Buran, Attitudes of health professionals toward elderly patients during the COVID-19 pandemic, *Aging Clin. Exp. Res.* (2022) 1–10, <https://doi.org/10.1007/s40520-022-02209-6>. Epub ahead of print.

Céline Demonceau^a, Charlotte Beaudart^a, Jean-Yves Reginster^a,
Nicola Veronese^b, Olivier Bruyère^{a,*}
^a University of Liege, Belgium
^b University of Palermo, Italy

* Corresponding author at: University of Liege, 4000, Belgium.
E-mail address: olivier.bruyere@uliege.be (O. Bruyère).